

HEAD LICE TREATMENT



Overview

Treatment for head lice is recommended for persons diagnosed with an active infestation. All household members and other close contacts should be checked; those persons with evidence of an active infestation should be treated. Some experts believe prophylactic treatment is prudent for persons who share the same bed with actively-infested individuals. All infested persons (household members and close contacts) and their bedmates should be treated at the same time.

Some pediculicides (medicines that kill lice) have an ovicidal effect (they kill eggs). For pediculicides that are only weakly ovicidal or not ovicidal, routine retreatment is recommended. For those that are more strongly ovicidal, retreatment is recommended only if live (crawling) lice are still present several days after treatment (see recommendation for each medication). To be most effective, retreatment should occur after all eggs have hatched but before new eggs are produced.

When treating head lice, supplemental measures can be combined with recommended medicine (pharmacologic treatment); however, such additional (non-pharmacologic) measures generally are not required to eliminate a head lice infestation. For example, hats, scarves, pillow cases, bedding, clothing and towels worn or used by the infested person in the two-day period just before treatment is started can be machine washed and dried using the hot water and hot air cycles, because lice and eggs are killed by exposure for five minutes to temperatures greater than 53.5°C (128.3°F). Items that cannot be laundered may be dry-cleaned or sealed in a plastic bag for two weeks. Items such as hats, grooming aids and towels that come in contact with the hair of an infested person should not be shared. Vacuuming furniture and floors can remove an infested person's hairs that might have viable nits attached.

Treatment of the infested person(s) requires using an over-the-counter (OTC) or prescription medication. Follow these treatment steps:

1. Before applying treatment, it may be helpful to remove clothing that can become wet or stained during treatment.
2. Apply lice medicine, also called pediculicide, according to the instructions contained in the box or printed on the label. If the infested person has very long hair (longer than shoulder length), it may be necessary to use a second bottle. Pay special attention to instructions on the label or in the box regarding how long the medication should be left on the hair and how it should be washed out of the hair.
3. Have the infested person put on clean clothing after treatment.
4. If a few live lice are still found eight to 12 hours after treatment, but are moving more slowly than before, do not retreat. The medicine may take longer to kill all the lice. Comb dead and any remaining live lice out of the hair using a fine-toothed nit comb.
5. If, after eight to 12 hours of treatment, no dead lice are found and lice seem as active as before, the medicine may not be working. Do not retreat until

speaking with your health care provider; a different pediculicide may be necessary. If your health care provider recommends a different pediculicide, carefully follow the treatment instructions contained in the box or printed on the label.

6. Nit (head lice egg) combs, often found in lice medicine packages, should be used to comb nits and lice from the hair shaft. Many flea combs made for cats and dogs are also effective.
7. After each treatment, checking the hair and combing with a nit comb to remove nits and lice every two to three days may decrease the chance of self-reinfestation. Continue to check for two to three weeks to be sure all lice and nits are gone. Nit removal is not needed when treating with spinosad topical suspension.
8. Retreatment is meant to kill any surviving hatched lice before they produce new eggs. For some drugs, re-treatment is recommended routinely about a week after the first treatment (seven to nine days, depending on the drug) and for others only if crawling lice are seen during this period. Retreatment with lindane shampoo is not recommended.

WARNING: Do not use a combination shampoo/conditioner or conditioner before using lice medicine. Do not re-wash the hair for one to two days after the lice medicine is removed. These measures may reduce the effectiveness of the lice medications.

Supplemental Measures: Head lice do not survive long if they fall off a person and cannot feed. You don't need to spend a lot of time or money on housecleaning activities. Follow these steps to help avoid re-infestation by lice that have recently fallen off the hair or crawled onto clothing or furniture.

1. Machine wash and dry clothing, bed linens and other items that the infested person wore or used during the two days before treatment using the hot water (130°F) laundry cycle and the high heat drying cycle. Clothing and items that are not washable can be dry-cleaned or sealed in a plastic bag and stored for two weeks.
2. Soak combs and brushes in hot water (at least 130°F) for five to 10 minutes.
3. Vacuum the floor and furniture, particularly where the infested person sat or lay. However, the risk of getting infested by a louse that has fallen onto a rug or carpet or furniture is very low. Head lice survive less than one to two days if they fall off a person and cannot feed; nits cannot hatch and usually die within a week if they are not kept at the same temperature as that found close to the human scalp. Spending much time and money on housecleaning activities is not necessary to avoid reinfestation by lice or nits that may have fallen off the head or crawled onto furniture or clothing.
4. Do not use fumigant sprays; they can be toxic if inhaled or absorbed through the skin.

Over-the-Counter Medications

Many head lice medications are available "over-the-counter" without a prescription at a local drug store or pharmacy. Each over-the-counter product approved by the

FDA for the treatment of head lice contains one of the following active ingredients. If crawling lice are still seen after a full course of treatment, contact your health care provider.

- Pyrethrins combined with piperonyl butoxide
Brand name products: A-200,* Pronto,* R&C,* Rid,* Triple X,* Licide*

Pyrethrins are naturally occurring pyrethroid extracts from the chrysanthemum flower. Pyrethrins are safe and effective when used as directed. Pyrethrins can only kill live lice, not unhatched eggs (nits). A second treatment is recommended nine to 10 days after the first treatment to kill any newly hatched lice before they can produce new eggs. Pyrethrins generally should not be used by persons who are allergic to chrysanthemums or ragweed. Pyrethrin is approved for use on children 2 years of age and older.

- Permethrin lotion, 1%;
Brand name product: Nix*

Permethrin is a synthetic pyrethroid similar to naturally occurring pyrethrins. Permethrin lotion 1% is approved by the FDA for the treatment of head lice. Permethrin is safe and effective when used as directed. Permethrin kills live lice but not unhatched eggs. Permethrin may continue to kill newly hatched lice for several days after treatment. A second treatment often is necessary on day nine to kill any newly hatched lice before they can produce new eggs. Permethrin is approved for use on children 2 months of age and older.

Prescription Medications

The following medications, in alphabetical order, approved by the U.S. Food and Drug Administration (FDA) for the treatment of head lice, are available only by prescription. If crawling lice are still seen after a full course of treatment, contact your health care provider.

- Benzyl alcohol lotion, 5%;
Brand name product: Ulesfia lotion*

Benzyl alcohol is an aromatic alcohol. Benzyl alcohol lotion, 5% has been approved by the FDA for the treatment of head lice and is considered safe and effective when used as directed. It kills lice, but it is not ovicidal (i.e., does not kill lice eggs). A second treatment is needed nine days after the first treatment to kill any newly hatched lice before they can produce new eggs. Benzyl alcohol lotion is intended for use on persons who are 6 months of age and older, and its safety for persons aged more 60 years has not been established. It can be irritating to the skin.

- Ivermectin lotion, 0.5%;
Brand name product: Sklice*

Ivermectin lotion, 0.5% was approved by the FDA in 2012 for treatment of head lice in persons 6 months of age and older. It is not ovicidal but appears to prevent nymphs (newly hatched lice) from surviving. It is effective in most patients when given as a single application on dry hair without nit combing. It should not be used for retreatment without talking to a health care provider.

Given as a tablet in mass drug administrations, oral ivermectin has been used extensively and safely for over two decades in many countries to treat filarial worm infections. Although not FDA-approved for the treatment of lice, ivermectin tablets given in a single oral dose of 200 micrograms/kg repeated in 10 days or 400 micrograms/kg repeated in seven days has been shown effective against head lice. It should not be used in children weighing less than 15 kg or in pregnant women.

- Malathion lotion, 0.5%;
Brand name product: Ovide*

Malathion is an organophosphate. The formulation of malathion approved in the United States for the treatment of head lice is a lotion that is safe and effective when used as directed. Malathion is pediculicidal (kills live lice) and ovicidal (kills lice eggs). A single application is adequate for most patients. However, a second treatment is recommended if live lice still are present seven to nine days after treatment. Malathion is intended for use on persons 6 years of age and older. Malathion can be irritating to the skin. Malathion lotion is flammable, so hair should be allowed to dry naturally; do not use electrical heat sources, including hair dryers, curlers, and curling or flat irons, or smoke when applying malathion lotion and while the hair is wet.

For more information on malathion, see http://www.cdc.gov/parasites/lice/head/gen_info/faqs_malathion.html

- Spinosad 0.9% topical suspension;
Brand name product: Natroba*

Spinosad is derived from soil bacteria. Spinosad topical suspension, 0.9% was approved by the FDA in 2011. Since it kills live lice as well as unhatched eggs, retreatment is usually not needed. Nit combing is not required. Spinosad topical suspension is approved for the treatment of children 6 months of age and older. It is safe and effective when used as directed. Repeat treatment should be given only if live (crawling) lice are seen seven days after the first treatment.

Second-Line Treatment

- Lindane shampoo, 1%;
Brand name products: None available

Lindane is an organochloride. The American Academy of Pediatrics (AAP) no longer recommends it as a pediculicide. Although lindane shampoo 1% is approved by the FDA for the treatment of head lice, it is not recommended as a first-line treatment. Overuse, misuse or accidentally swallowing lindane can be toxic to the brain and other parts of the nervous system; its use should be restricted to patients for whom prior treatments have failed or who cannot tolerate other medications that pose less risk. Lindane should not be used to treat premature infants, persons with HIV or a seizure disorder, women who are pregnant or breastfeeding, persons who have very irritated skin or sores where the lindane will be applied, infants, children, older residents, and persons who weigh less than 110 pounds. Retreatment should be avoided.

When treating head lice:

- Do not use extra amounts of any lice medication unless instructed to do so by your physician or pharmacist. The drugs used to treat lice are insecticides and can be dangerous if they are misused or overused.
- All the medications listed above should be kept out of the eyes. If they get onto the eyes, they should be immediately flushed away.
- Do not treat an infested person more than two to three times with the same medication if it does not seem to be working. This may be caused by using the medicine incorrectly or by resistance to the medicine. Always seek the advice of your health care provider if this should happen. He/she may recommend an alternative medication.
- Do not use different head lice drugs at the same time unless instructed to do so by your physician or pharmacist.

* These are commercial products and are noted for the convenience of the reader and do not imply any endorsement nor recommendation by the Pennsylvania Department of Health.

Additional Information

Centers for Disease Control and Prevention Head Lice Treatment:

<http://www.cdc.gov/parasites/lice/head/treatment.html>

Mayo Clinic Head Lice Treatments and Drugs: <http://www.mayoclinic.org/diseases-conditions/head-lice/basics/treatment/con-20030792>

National Pediculosis Association Treatments:

<http://www.headlice.org/faq/treatments/index.htm>

Last reviewed/updated: June 20, 2016